

Table 21.7
Cancer of the Ovary (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2006-2010</u>						
Age at Diagnosis/Death:						
All ages	12.5	13.3	9.7	8.1	8.4	6.7
Under 65	7.6	8.0	5.4	3.2	3.3	2.7
65 and over	46.6	49.5	39.6	42.0	43.7	34.3
All ages (IARC world std) ^c	8.9	9.4	6.8	5.1	5.3	4.3
<u>Age-Specific Rates, 2006-2010</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	0.3	0.2	-	-	-	-
10-14	0.8	0.8	-	-	-	-
15-19	1.3	1.4	1.0	0.1	0.0	-
20-24	1.7	1.8	1.4	0.2	0.2	-
25-29	2.3	2.4	2.0	0.3	0.3	0.4
30-34	3.3	3.4	2.7	0.6	0.6	0.7
35-39	4.8	5.0	3.6	1.0	1.1	1.1
40-44	8.9	9.4	5.5	2.4	2.5	2.2
45-49	14.2	14.9	9.3	4.9	5.2	4.2
50-54	20.7	21.6	14.0	9.0	9.4	7.4
55-59	26.1	28.1	18.0	14.0	14.8	11.4
60-64	32.5	34.7	25.3	21.0	22.0	17.4
65-69	39.3	42.3	33.1	29.0	30.2	25.4
70-74	45.6	48.7	38.4	38.0	39.8	31.4
75-79	49.4	52.8	40.4	46.0	47.7	37.8
80-84	53.0	55.4	46.6	54.8	56.8	42.8
85+	52.5	54.4	47.2	57.6	59.3	44.3

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473. SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.